IN THE CLAIMS

Please amend the claims as follows:

Claim 1 (Previously Presented): A herbicidal composition which comprises i) an isoxazoline derivative represented by the following general formula (I) or a salt

$$R^{1} \xrightarrow{R^{2}} R^{3}$$

$$R^{4} \xrightarrow{R^{6}} S(0)_{n} \xrightarrow{R^{6}} Y$$

$$R^{5} \xrightarrow{R^{5}} Y$$

$$R^{1} \xrightarrow{R^{6}} Y$$

wherein R¹ and R² are independently a hydrogen atom, a C1 to C10 alkyl group, a C3 to C8 cycloalkyl group or a C3 to C8 cycloalkyl C1 to C3 alkyl group; or R¹ and R² may be bonded to each other to form a C3 to C7 spiro ring together with the carbon atoms to which they bond;

R³ and R⁴ are independently a hydrogen atom, a C1 to C10 alkyl group or a C3 to C8 cycloalkyl group; or R³ and R⁴ may be bonded to each other to form a C3 to C7 spiro ring together with the carbon atoms to which they bond; or R¹, R², R³ and R⁴ may form a 5- to 8-membered ring together with the carbon atoms to which they bond;

 R^5 and R^6 are independently a hydrogen atom or a C1 to C10 alkyl group;

Y is a 5- to 6-membered aromatic heterocyclic group or condensed aromatic heterocyclic group having one or more hetero atoms selected from a nitrogen atom, an oxygen atom and a sulfur atom; the heterocyclic group may be substituted with 0 to 6 same or different groups selected from the following substituent group α ; when the heterocyclic group is substituted at the two adjacent positions with two alkyl groups, two alkoxy groups, an alkyl group and an alkoxy group, an alkyl group and an alkylthio group, an alkyl group and an alkylsulfonyl group, an alkyl group and a monoalkylamino group, or an alkyl group and a dialkylamino group, all selected from the substituent group α , the two groups may form,

together with the atoms to which they bond, a 5- to 8-membered ring which may be substituted with 1 to 4 halogen atoms; the hetero atom of the heterocyclic group, when it is a nitrogen atom, may be oxidized to become N-oxide;

n is an integer of 0 to 2;

wherein said substituent group α is selected from the group consisting of hydroxyl group; thiol group; halogen atoms; C1 to C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the following substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group selected from the following substituent group γ; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkyloxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylthio groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl groups; C2 to C6 alkenyloxy groups; C2 to C6 alkynyl groups; C2 to C6 alkynyloxy groups; C1 to C10 alkylsulfinyl groups; C1 to C10 alkylsulfinyl groups each mono-substituted with a group selected from the substituent group γ ; C1 to C10 alkylsulfonyl groups; C1 to C10 alkylsulfonyl groups each mono-substituted with a group selected from the substituent group γ; C1 to C4 haloalkylsulfinyl groups; C1 to C10 alkylsulfonyloxy groups each mono-substituted with a group selected from the substituent group γ; C1 to C4 haloalkylsulfonyl groups; C1 to C10 alkylsulfonyloxy groups; C1 to C4 haloalkylsulfonyloxy groups; optionally substituted phenyl group; optionally substituted phenoxy group; optionally substituted phenylthio group; optionally substituted aromatic heterocyclic groups; optionally substituted aromatic heterocyclic oxy groups; optionally substituted aromatic heterocyclic thio groups; optionally substituted phenylsulfinyl groups; optionally substituted phenylsulfonyl groups; optionally substituted aromatic heterocyclic

sulfonyl groups; optionally substituted phenylsulfonyloxy groups; acyl groups; C1 to C4 haloalkylcarbonyl groups; optionally substituted benzylcarbonyl group; optionally substituted benzyloxycarbonyl group; carboxyl group; C1 to C10 alkoxycarbonyl groups; optionally substituted benzyloxycarbonyl group; optionally substituted phenoxycarbonyl group; cyano group; carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group); C1 to C6 acyloxy groups; C1 to C4 haloalkylcarbonyloxy groups; optionally substituted benzylcarbonyloxy group; optionally substituted benzoloxy group; nitro group; and amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, C1 to C6 acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl group, optionally substituted benzylcarbonyl group, optionally group, optionally substituted

wherein said substituent group β is selected from the group consisting of hydroxyl group; C3 to C8 cycloalkyl groups (which may be substituted with halogen atom or alkyl group); C1 to C10 alkoxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylsulfonyl groups; C1 to C10 alkoxycarbonyl groups; C2 to C6 haloalkenyl groups; amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups, C1 to C10 alkylsulfonyl groups and C1 to C4 haloalkylsulfonyl groups); carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups); C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups; C1 to C10 alkoxyimino groups; cyano group; optionally substituted phenyl group; and optionally substituted phenoxy group;

wherein said substituent group γ is selected from the group consisting of

C1 to C10 alkoxycarbonyl groups; optionally substituted phenyl group; optionally substituted aromatic heterocyclic groups; cyano group; and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups); and

ii) at least one compound selected from the group consisting of

atrazine, simazine, cyanazine, isoxaflutole, mesotrione, flumetsulam, imazethapyr, imazapyr, dicamba, clopyralid, prosulfuron, halosulfuron-methyl, rimsulfuron, bentazone, carfentrazone-ethyl, metribuzin, thifensulfuron-methyl, nicosulfuron, primisulfuron, cloransulam-methyl, glufosinate, glyphosate, glyphosate-trimesium, pendimethalin, linuron, prometryn, diflufenican, flumioxazin, and metolachlor.

Claim 2 (Withdrawn): The herbicidal composition according to Claim 1, wherein the isoxazoline derivative of the formula (I) or a salt thereof has a substituent selected from the substituent group α on the heterocycle which may be substituted with 0 to 6 same or different groups, including hydroxyl group; halogen atoms; C1 to C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkyloxy groups; C3 to C8 cycloalkyloxy groups; C1 to C10 alkylthio groups; C1 to C3 alkyloxy groups; C1 to C10 alkylthio groups; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl groups; C2 to C6 alkenyloxy groups; C2 to C6 alkenyloxy groups; C1 to C10 alkylsulfonyl groups; C1 to C4 haloalkylsulfonyl groups; optionally substituted phenoxy group; optionally substituted phenylthio group; optionally substituted aromatic heterocyclic groups; optionally substituted aromatic heterocyclic thio groups; optionally substituted aromatic heterocyclic thio groups; optionally

substituted phenylsulfonyl groups; optionally substituted aromatic heterocyclic sulfonyl groups; C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups; optionally substituted benzylcarbonyl group; optionally substituted benzoyl group; carboxyl group; C1 to C10 alkoxycarbonyl groups; cyano group; carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group); nitro group; and amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, C1 to C6 acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group, and optionally substituted phenylsulfonyl group); when the heterocyclic group is substituted at the two adjacent positions with two alkyl groups, two alkoxy groups, an alkyl group and an alkoxy group, an alkyl group and an alkylthio group, an alkyl group and an alkylsulfonyl group, an alkyl group and a monoalkylamino group, or an alkyl group and a dialkylamino group, the two groups may form, together with the atoms to which they bond, a 5- to 8-membered ring which may be substituted with 1 to 4 halogen atoms.

Claim 3 (Withdrawn): The herbicidal composition according to Claim 2, wherein the isoxazoline derivative of the formula (I) or a salt thereof has a substituent selected from the substituent group α on the heterocycle which may be substituted with 0 to 6 same or different groups, including halogen atoms; C1 to C10 alkyl groups; C1 to C4 haloalkyl groups; C1 to C10 alkoxy C1 to C3 alkyl groups; C3 to C8 cycloalkyl groups (which may be substituted with halogen atom or alkyl group); C1 to C10 alkoxy groups; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; optionally substituted phenoxy group; C1 to

C10 alkylthio groups; C1 to C10 alkylsulfonyl groups; acyl groups; C1 to C4 haloalkylcarbonyl groups; C1 to C10 alkoxycarbonyl groups; cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups).

Claim 4 (Previously Presented): The herbicidal composition according to Claim 1, wherein R¹ and R² may be the same or different and are each a methyl group or an ethyl group; and R³, R⁴, R⁵ and R⁶ are each a hydrogen atom.

Claim 5 (Previously Presented): The herbicidal composition according to Claim 1, wherein Y is a 5- or 6-membered aromatic heterocyclic group having a hetero atom selected from a nitrogen atom, an oxygen atom and a sulfur atom.

Claim 6 (Original): The herbicidal composition according to Claim 5, wherein Y is a thienyl group, a pyrazolyl group, an isoxazolyl group, an isothiazolyl group, a pyridyl group or a pyrimidinyl group.

Claim 7 (Original): The herbicidal composition according to Claim 6, wherein Y is a thiophen-3-yl group, a pyrazol-4-yl group, a pyrazol-5-yl group, an isoxazol-4-yl group, an isothiazol-4-yl group, a pyridyn-3-yl group or a pyrimidin-5-yl group.

Claim 8 (Original): The herbicidal composition according to Claim 7, wherein Y is a thiophen-3-yl group and the thiophene ring is substituted with the substituent group α at the 2- and 4-positions.

Claim 9 (Original): The herbicidal composition according to Claim 7, wherein Y is a pyrazol-4-yl group and the pyrazole ring is substituted at the 3- and 5-positions with the substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent group β , a C1 to C4 haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl group, a C1 to C4 haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxycarbonyl group, an optionally substituted benzyloxycarbonyl group, an optionally substituted phenoxycarbonyl group, a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group), or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, an optionally substituted phenyl group, acyl groups, C1 to C4 haloalkylcarbonyl groups, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, an optionally substituted benzylsulfonyl group and an optionally substituted phenylsulfonyl group).

Claim 10 (Original): The herbicidal composition according to Claim 7, wherein Y is a pyrazol-5-yl group and the pyrazole ring is substituted at the 4-position with the substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent group β , a C1 to C4

haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl group, a C1 to C4 haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxycarbonyl group, an optionally substituted benzyloxycarbonyl group, an optionally substituted phenoxycarbonyl group, a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and an optionally substituted phenyl group), or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, an optionally substituted phenyl group, acyl groups, C1 to C4 haloalkylcarbonyl groups, an optionally substituted benzylcarbonyl group, an optionally substituted benzyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, an optionally substituted benzylsulfonyl group and an optionally substituted phenylsulfonyl group).

Claim 11 (Original): The herbicidal composition according to Claim 7, wherein Y is an isoxazol-4-yl group and the isoxazole ring is substituted with the substituent group α at the 3- and 5-positions.

Claim 12 (Original): The herbicidal composition according to Claim 7, wherein Y is an isothiazol-4-yl group and the isothiazole ring is substituted with the substituent group α at the 3- and 5-positions.

Claim 13 (Original): The herbicidal composition according to Claim 7, wherein Y is a pyridin-3-yl group and the pyridine ring is substituted with the substituent group α at the 2-and 4-positions.

Claim 14 (Original): The herbicidal composition according to Claim 7, wherein Y is a pyrimidin-5-yl group and the pyrimidine ring is substituted with the substituent group α at the 4- and 6-positions.

Claim 15 (Previously Presented): The herbicidal composition according to Claim 1, wherein n is an integer of 2.

Claim 16 (Currently Amended): The herbicidal composition according to Claim 1, wherein the compound of Group A ii) is at least one compound selected from the group consisting of atrazine, cyanazine, simazine and prometryn.

Claim 17 (Currently Amended): The herbicidal composition according to Claim 1, wherein the compound of Group A ii) is at least one compound selected from the group consisting of glyphosate, glufosinate, linuron and flumetsulam.

Claim 18 (Currently Amended): A herbicidal composition which comprises i) the isoxazoline derivative or a salt thereof is a compound as defined in Claim 9 and [[ii)]] the compound of Group A ii) is at least one compound selected from the group consisting of atrazine, cyanazine, simazine, prometryn, glyphosate, glufosinate, linuron, flumetsulam, metribuzin, isoxaflutole, mesotrione, diflufenican, pendimethalin and flumioxazin.

Claim 19 (Currently Amended): A herbicidal composition which comprises i) the isoxazoline derivative or a salt thereof is a compound as defined in Claim 9 and [[ii)]] the compound of Group A ii) is at least one compound selected from the group consisting of atrazine, cyanazine, simazine and prometryn.

Claim 20 (Currently Amended): A herbicidal composition which comprises i) the isoxazoline derivative or a salt thereof is a compound as defined in Claim 9 and [[ii)]] the compound of Group A ii) is at least one compound selected from the group consisting of glyphosate, glufosinate, linuron and flumetsulam.

Claim 21 (Currently Amended): The herbicidal composition according to Claim 1, wherein ii) at least one the compound of Group A ii) is comprised in an amount of from 0.001 to 100 parts by weight to 1 part by weight of i) an isoxazoline derivative represented by the Formula (I) or a salt thereof.

Claim 22 (Currently Amended): The herbicidal composition according to Claim 1, which is used as an agrochemical product comprising i) an isoxazoline derivative of the Formula (I) or a salt thereof and ii) at least one the compound of Group A ii) in a total amount of from 0.5 to 90 wt%.

11